



Grey Kitty Lawn Care Program

Late March/April	Apply pre-emergent to prevent weed germination. Do not apply if over-seeding; Apply Granular Revive to benefit from spring moisture.
April/May	Core aerate and apply Natural Guard® Granular Humic; Top dress with ¼" compost or EKO Lawn Top Dressing; Seed or sod if necessary.
May/June	Apply a slow-release fertilizer. Water-in.
July/August	Apply Revive® for heat stress and improved water penetration
Late Aug./Early Sept.	Apply pre-emergent to prevent weed germination. Do not apply if over-seeding.
September	Core aerate and apply Natural Guard® Granular Humic; Top dress with ¼" compost or EKO Lawn Top Dressing; Seed or sod if necessary.
Oct/November	Apply a slow-release Winterizer , water-in.

Slow-Release Fertilizers: Richlawn **Pro-Rich**, Richlawn **Organic 100** and **Colorado's Own Lawn Food**

Revive® improves water penetration, adds nutrients, and promotes greener turf. Apply any time during growing season if needed; Available in both granular and liquid forms.

Richlawn® Iron-Rich helps lawns that are healthy, but are pale green; only apply once during growing season.

F-Stop is a fungicide that is helpful for several common lawn diseases. It should be applied (if needed) in spring to **prevent** fungal disease; later application will help disease from spreading.

EKO Lawn Top Dressing can be applied ¼" thick to thinning and stressed turf areas. This sifted compost will enrich soil, promote root growth, and improve turf health.

Natural Guard Granular Humic can be applied anytime to improve soil quality.

****Helpful Hints:**

Read all instructions on package labels before applying; over-applying or mixing incorrectly can cause damage.

Don't fertilize too close to seeding; only after 2nd mowing can seedlings handle fertilizer.

If weeds occur, remove manually or spot spray with an herbicide; apply in the evening when temperature is between 65 and 85 with no wind.

Mow grass 3" tall to block weeds, grow deeper roots, and retain water.

Water 1 ½" per week (broken into 3- ½" morning cycles). Increase up to 2-3" during hottest part of the summer. **Never water an established lawn more than every 2nd day; Adjust according to rainfall and temperature.**